

Scientific note

urn:lsid:zoobank.org:pub:0D573D65-6DD1-4B3C-9ED9-D44F48638C54

First record of males of the invasive eucalyptus pest species *Leptocybe invasa* Fisher & LaSalle, 2004 (Hymenoptera: Eulophidae: Tetrastichinae) from South America

Daniel Alejandro Aquino^{1,*}, Carmen Marcela Hernández² & Andrea Verónica Andorno³

¹Centro de Estudios Parasitológicos y de Vectores (CEPAVE, CONICET-UNLP), Boulevard 120 entre 60 y 64,
AR-1900 La Plata, Argentina

^{2,3}Instituto Nacional de Tecnología Agropecuaria (INTA). Instituto de Microbiología y Zoología Agrícola, AR-Argentina

*Corresponding author: Email: daquino@fcnym.unlp.edu.ar

¹urn:lsid:zoobank.org:author:670E065F-04CF-4247-8515-AC41D151017A

²urn:lsid:zoobank.org:author:778230C8-8EA2-4C21-892B-ACB488810D25

³urn:lsid:zoobank.org:author:47B448E3-6B06-4343-BB2F-8E87D4E17750

Abstract. The “blue gum chalcid”, *Leptocybe invasa* is a pest of eucalyptus worldwide. It has an Australian origin and has expanded into Asia, Europe, Africa and America. *L. invasa* females were reported from South America in the last decade. Thelytokous parthenogenesis is the most common reproductive mechanism of this pest. However, male adults have been reported from Asia and Southeastern Europe. In this work, *L. invasa* males are reported for the first time from South America (Argentina) and information on morphological characters of males is provided. The importance of this discovery is highlighted based on recent studies that suggested the existence of two cryptic *Leptocybe* species in invasive populations.

Key words. Blue gum chalcid, gall, forest, chalcidoidea.

The “blue gum chalcid”, *Leptocybe invasa* Fisher & LaSalle, 2004 (Hymenoptera: Eulophidae: Tetrastichinae) is a global pest in *Eucalyptus* plantations. The wasp lays eggs in plant tissues causing the formation of galls on the midribs, petioles and stems of young leaves. Severely infested trees have a gnarled appearance, show stunted growth, and in young trees, this pest can cause significant injury (Mendel et al. 2004; Nyeko, 2005).

The rapid colonization after *L. invasa* spread into a novel environment can be attributed to the absence of natural enemies, the presence of large amounts of suitable host plants, the resistance to low temperatures of adults and the occurrence of two reproductive modes (Zheng et al. 2014b). This pest displays thelytokous reproduction but since the discovery of males, sexual reproduction has been considered a second reproductive strategy. Nevertheless, factors influencing reproductive strategies and the relation between reproductive strategy and population expansion are unknown (Zheng et al. 2014a). In the current study, we describe male specimens of *L. invasa* from Argentina and report the sex ratio in eucalyptus plantations in the country.

In 2017 and 2018, a survey of *L. invasa* populations and natural enemies was conducted in infested eucalyptus plantations in an extensive productive area in Buenos Aires and the Mesopotamia region (Entre Ríos, Corrientes and Misiones). Almost fifty samples of 20–30 cm

long, *L. invasa* gall infested branches were collected. The plant material was kept in glass containers with voile lid and absorbent paper under controlled laboratory conditions (T: 25 ± 2°C, HR: 50–70% and natural lighting). The insects that emerged from each sample were kept in 70% ethanol and were identified by the first author. The specimens were card- and slide-mounted following Noyes (1990) and were compared with the descriptions of the males in Doğanlar (2005), Chen et al. (2009) and Zheng et al. (2014b). Voucher specimens were deposited at the División Entomología of Museo de La Plata. Sex ratios were calculated as $\Sigma \text{♀} / \Sigma (\text{♀} + \text{♂})$. Measurements were taken with a Leitz Wetzlar SM/LUX; specimens were photographed using a Leica DFC295 digital camera attached to a stereomicroscope Biotraza. Series of partially focused digital images were stacked using the Helicon Focus software (Version 6.8.0 Pro) by ©Helicon Soft Ltd., 2000. Photoshop CC2018 was used to produce final images with enhanced quality.

Material examined. Argentina, Entre Ríos, Concordia en 8.iii.2018 and 5.x.2018 (sex ratio 3m:104f (0,972) and 17m:1290f (0.987)); Argentina, Corrientes, Alvear 28.ix.2018 (sex ratio 1m:24f (0,960)).

According to Doğanlar (2005), the male is similar to the female described by Fisher & LaSalle (2004), except:



Figs. 1–4. *Leptocybe invasa* male from Argentina. 1. lateral habitus; 2. antenna; 3. genitalia; 4. forewing.

head and mesosoma brown with distinct blue to green metallic shine (Fig. 1); ocellar triangle and mouth margin light brown; eyes and ocellus red; malar sulcus dark brown; metasoma brown; legs almost white, except mid and hind coxae same color of the body and hind femur light brown. Antenna with scape and pedicel light brown darkened dorsally and apically; funicle and clava light brown to almost white. Wings hyaline, veins light brown (Fig. 4). Genitalia (Fig. 3) with digitus having one claw. The specimens collected in this study are similar to those described by Doğanlar (2005) except for the following measurements and ratios: Head 1.28 times as broad as high. Antenna (Fig. 2) with scape 3.13 times as long as broad, ventral plaque 0.18–0.24x length of scape; pedicel 1.9–2.8x as long as broad, longer than anelli plus F1; F1 shorter than F2 (in figure 1b in Doğanlar (2005) it is observable that F1 is shorter than F2 although he mentions

that “F1 is 1.46x longer than F2”), about 0.75x length of F2, as long as broad to 0.7 times as long as broad, F2 to F4 subequal in length, each slightly longer than broad; clava about 2.75 times as long as broad, i.e., slightly longer than that described by Doğanlar (2005), with C1–C3 decreasing in length, except in one specimen in which C1 is shorter than C2 and C3.

Leptocybe invasa males were reported from Turkey, China, India, Taiwan and Thailand, and populations of this pest around the world are known to have different sex ratios over their geographic distribution (Nugnes et al. 2015). Males are rare in Turkey (sex ratio: 0,992, n=125) (Doğanlar 2005) and India (sex ratio: 0,992, n=141) (Akhtar et al. 2012). In China a low proportion of males was reported from Fujian, Guangdong, Hainan, Guangxi, Jiangxi and Sichuan (sex ratio $\geq 0,95$), but a high proportion of males was reported from other regions (males proportion ranging from 18–48%) and in Thailand (sex ratio: 0,663) (Liang et al. 2010; Santongprow et al. 2011; Zheng et al. 2018).

In this work, in only two out of twenty sampling sites males were found (in Entre Ríos and in Corrientes) and a clearly female-biased sex ratio was found, with values $\geq 0,96$.

While *L. invasa* females were recorded from several countries in America since 2007 (Brazil (2007), Argentina (2009), Paraguay (2012), Uruguay (2013), Chile (2014), Mexico (2014) and USA (2008) (Costa et al. 2008; Wiley and Skelley, 2008; Aquino et al. 2011; SAG 2014; Benítez et al. 2014; Vanegas et al. 2015; Jorge et al. 2016)), this is the first report of *L. invasa* males from America. Recent molecular analyses suggested that *L. invasa* is, in fact, a complex of at least two cryptic species involved in the rapid and efficient spread of the wasp (Nugnes et al. 2015; Dittrich-Schröder et al. 2018). Nugnes et al. (2015) proposed that there is a link between the presence of males in the invasive range and lineage identity. Further molecular analyses are necessary to understand the invasion pathways and to characterize the populations of *Leptocybe* species in Argentina and other countries in America.

Acknowledgements. The authors thank Sergio Ramos, Cintia Meneses and Edgar Eskiviski for providing samples. We gratefully acknowledge to CONICET and the financial support from PICT-2016-0834 (FONCYT, MINCYT, Argentina).

REFERENCES

- Akhtar MS, Patankar NV, Gaur A (2012) Observations on the biology and male of eucalyptus gall wasp *Leptocybe invasa* Fisher & LaSalle (Hymenoptera: Eulophidae). Indian Journal of Entomology 74: 173–175
- Aquino DA, Botto EN, Loíacono MS, Pathauer P (2011) Avispa de la agalla del Eucalipto, *Leptocybe invasa* Fisher & LaSalle (Hymenoptera: Eulophidae: Tetrastichinae) en Argentina. Revista de Investigaciones Agropecuarias 37 (2): 159–164

- Benítez EA, Costa VA, de Moraes GJ, Godziewsky D (2014) First record of *Leptocybe invasa* Fisher & LaSalle (Hymenoptera: Eulophidae) and *Rhombacus eucalypti* Ghosh & Chakrabarti (Acari: Eriophyidae) from Paraguay. Boletín del Museo Nacional de Historia Natural de Paraguay 18: 129–132
- Chen HY, Yao JM, Xu ZF (2009) First description of the male of *Leptocybe invasa* Fisher & LaSalle (Hymenoptera: Eulophidae) from China. Journal of Environmental Entomology 31: 285–287 (in Chinese with English abstract)
- Costa VA, Filho EB, Wilcken CF, Stape JL, LaSalle J, Teixeira L (2008) *Eucalyptus* gall wasp, *Leptocybe invasa* Fisher & LaSalle (Hymenoptera: Eulophidae) in Brazil: New forest pest reaches the New World. Revista de Agricultura 83: 136–139
- Dittrich-Schröder G, Hoareau TB, Hurley BP, Wingfield MJ, Lawson S, Nahrung HF, Slippers B (2018) Population genetic analyses of complex global insect invasions in managed landscapes: a *Leptocybe invasa* (Hymenoptera) case study. Biological Invasions 20: 2395
- Doğanlar O (2005) Occurrence of *Leptocybe invasa* Fisher & LaSalle, 2004 (Hymenoptera: Chalcidoidea: Eulophidae) on *Eucalyptus camaldulensis* in Turkey, with description of the male sex. Zoology in the Middle East 35: 112–114
- Jorge C, Martínez G, Gómez D, Bollazzi M (2016) First record of the eucalypt gall-wasp *Leptocybe invasa* (Hymenoptera: Eulophidae) from Uruguay. BOSQUE 37(3): 631–636
- Kumari KN, Kulkarni H, Vastrad AS, Goud KB (2010) Biology of eucalyptus gall wasp, *Leptocybe invasa* Fisher & LaSalle (Hymenoptera: Eulophidae). Karnataka Journal of Agricultural Sciences 23: 211–212
- Liang YP, Li JY, Wen XJ, Li YX (2010) Observation on the male ratio of *Leptocybe invasa* adult. Forest Pest and Disease 29: 21–22 (in Chinese with English abstract).
- Mendel Z, Protasov A, Fisher N, LaSalle J (2004) Taxonomy and biology of *Leptocybe invasa* gen. & sp. n. (Hymenoptera: Eulophidae), an invasive gall inducer on eucalyptus. Australian Journal of Entomology 43: 101–113
- Nugnes F, Gebiola M, Monti MM, Gualtieri L, Giorgini M, Wang JG, Bernardo U (2015) Genetic diversity of the invasive gall wasp *Leptocybe invasa* (Hymenoptera: Eulophidae) and of its Rickettsia endosymbiont, and associated sex ratio differences. PLoS One 10(5): e0124660. <https://doi.org/10.1371/journal.pone.0124660>
- SAG (Servicio Agrícola y Ganadero, CL) (2014) *Leptocybe invasa* Fischer & LaSalle (Hymenoptera: Eulophidae) Microavispa formadora de agallas en eucalipto. Servicio Agrícola y Ganadero. <https://www.sag.gob.cl/ambitos-de-accion/leptocybe-invasa-fischer-lasalle-hymenoptera-eulophidae-microavispa-formadora-de>
- Sangtongpraow B, Charernsom K, Siripatanadilok S (2011) Longevity, fecundity and development time of Eucalyptus gall wasp, *Leptocybe invasa* Fisher and LaSalle (Hymenoptera: Eulophidae) in Kanchanaburi province, Thailand. Thai Journal of Agricultural Science 44: 155–163
- Tung GS, LaSalle J (2010) Pest alert – a newly discovered invasion of gall forming wasps, *Leptocybe invasa* Fisher & LaSalle, on eucalyptus trees in Taiwan. Formosan Entomology 30: 241–244
- Vanegas-Rico JM, Lomeli-Flores JR, Rodríguez-Leyva E, Jiménez-Quiroz E, Pujade-Villar J (2015) First record of eucalyptus gall wasp *Leptocybe invasa* (Hymenoptera: Eulophidae) in Mexico. Revista Mexicana de Biodiversidad 86(4): 1095–1098
- Wiley J, Skelley PA (2008) A eucalyptus pest, *Leptocybe invasa* Fisher & LaSalle (Hymenoptera: Eulophidae), genus and species new to Florida and North America. Pest Alert. Florida, USA: Florida Department of Agriculture and Consumer Services, Division of Plant Industry. http://www.freshfromflorida.com/content/download/68487/1614796/Pest_Alert_-_Leptocybe_invasa_Blue_Gum_Chalcid.pdf
- Zheng X, Huang Z, Li J, Yang Z, Lu W (2018) Reproductive Biology of *Leptocybe invasa* Fisher & LaSalle (Hymenoptera: Eulophidae). Neotropical Entomology 47(1): 19–25
- Zheng XL, Li J, Yang ZD, Xian ZH, Wei JG, Lei CL, Wang XP, Lu W (2014a) A review of invasive biology, prevalence and management of *Leptocybe invasa* Fisher & LaSalle (Hymenoptera: Eulophidae: Tetrastichinae). African Entomology 22: 68–79
- Zheng XL, Yang ZD, Li J, Xian ZH, Yang J, Liu JY, Su S, Wang XL, Lu W (2014b) Rapid identification of both sexes of *Leptocybe invasa* Fisher & LaSalle (Hymenoptera: Eulophidae: Tetrastichinae): a morphological perspective. African Entomology 22(3): 643–650